

Basic Information

Product Name	Anti-GAPDH Antibody	
Gene Name	GAPDH	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat, monkey, zebrafish, chicken	
Tested Application	WB, IHC, ICC/IF, FCM	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Observed MW	36 kDa	
Dilution Ratios	Western blot (WB): Immunohistochemistry (IHC): Immunocytochemistry/Immunofluorescence (ICC/IF): Flow Cytometry (Fixed): (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	1:5000-20000 1:50-400 1:50-400 1:50-200

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Reference

Anti-GAPDH Antibody被引用在115文献中。

Selected Validation Data




Figure 1. Western blot analysis of anti- GAPDH antibody (A00227-1). The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: HeLa whole cell lysates,

Lane 2: Caco-2 whole cell lysates,

Lane 3: CCRF-CEM whole cell lysates,

Lane 4: rat brain tissue lysates,

Lane 5: rat liver tissue lysates,

Lane 6: mouse brain tissue lysates,

Lane 7: mouse liver tissue lysates.

Use rabbit anti- GAPDH 1:1000, probed with a goat anti-rabbit IgG-
HRP secondary antibody. The signal is developed using an
Enhanced Chemiluminescent detection (ECL) kit (Catalog #
EK1002). A specific band was detected for GAPDH at approximately
36KD. The expected band size for GAPDH is at 36KD.




Figure 2. IHC analysis using anti- GAPDH antibody
(A00227-1).detected in paraffin-embedded section of human
Laryngeal squamous cell carcinoma tissue. Biotinylated goat anti-
rabbit IgG was used as secondary antibody. The tissue section was
developed using Streptavidin-Biotin-Complex (SABC) (Catalog #
SA1022) with DAB as the chromogen.