

## Basic Information

<b>Product Name</b>	Anti-(Mouse) IHH Antibody (N-term)
<b>Gene Name</b>	Ihh
<b>Source</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Species Reactivity</b>	mouse
<b>Tested Application</b>	WB, FCM, IHC
<b>Contents</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
<b>Immunogen</b>	This mouse Ihh antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 57-90 amino acids from the N-terminal region of mouse Ihh.
<b>Purification</b>	This antibody is purified through a protein A column, followed by peptide affinity purification

## Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

## Background Information

Intercellular signal essential for a variety of patterning events during development. Binds to the patched (PTC) receptor, which functions in association with smoothened (SMO), to activate the transcription of target genes. Implicated in endochondral ossification: may regulate the balance between growth and ossification of the developing bones. Induces the expression of parathyroid hormone-related protein (PTHrP).

## Selected Validation Data

Product datasheet

**Anti-(Mouse) IHH Antibody (N-term)**

**Catalog Number: M01448**


**BOSTER**

antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator,  
Wuhan East Lake High-tech Development Zone

**Web:** www.boster.com **Phone:** 027-67845390 **Email:** boster@boster.com



Anti-Ihh Antibody (N-term) at 1:2000 dilution + mouse kidney lysates

Lysates/proteins at 20 µg per lane.

Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution

Predicted band size : 45 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.