BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator, Wuhan East Lake High-tech Development Zone

antibody and ELISA

Web: www.boster.com Phone: 027-67845390 Email: boster@boster.com

Basic Information	
Product Name	Anti-Flag Tag Antibody
Source	Mouse
Clonality	Monoclonal
lsotype	lgG1
Species Reactivity	human, mouse, rat, All species
Tested Application	WB, ICC/IF, IP
Contents	Mouse IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.5mg/ml BSA.
Immunogen	A synthesized peptide
Concentration	500ug/ml
Purification	Affinity-chromatography
Dilution Ratios	Western blot (WB):1:3000-30000Immunocytochemistry/Immunofluorescence (ICC/IF):1:50-200ImmunoPrecipitation (IP):1:100

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Reference

Anti-Flag Tag Antibody被引用在7文献中。

Selected Validation Data



BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator, Wuhan East Lake High-tech Development Zone

Web: www.boster.com Phone: 027-67845390 Email: boster@boster.com

Western blot analysis of Flag-tagged protein expression in (1) 293T cell lysate; (2) 293T cell transfected with Flag-tagged protein lysate.



