

Basic Information

Product Name	Anti-COX2/Cyclooxygenase 2/PTGS2 Antibody	
Gene Name	PTGS2	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse	
Tested Application	WB, IHC, FCM, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E.coli-derived human COX2/Cyclooxygenase 2/PTGS2 recombinant protein (Position: A18-L604).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	75 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-400
	Flow Cytometry (Fixed):	1:50-200
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000
	(Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


Background Information

Prostaglandin-endoperoxide synthase 2,also known as cyclooxygenase-2 or COX-2,is an enzyme that in humans is encoded by the PTGS2 gene. It is mapped to 1q31.1. Prostaglandin-endoperoxide synthase (PTGS),also known as cyclooxygenase,is the key enzyme in prostaglandin biosynthesis,and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2,which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events,suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis.

Reference

Anti-COX2/Cyclooxygenase 2/PTGS2 Antibody被引用在11文献中。

Selected Validation Data



Western blot analysis of COX2/Cyclooxygenase 2/PTGS2 using anti-COX2/Cyclooxygenase 2/PTGS2 antibody (A00084-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.


Lane 1: human Hela whole cell lysates,

Lane 2: mouse RAW264.7(-LPS) whole cell lysates,

Lane 3: mouse RAW264.7(+LPS) whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-COX2/Cyclooxygenase 2/PTGS2 antigen affinity purified polyclonal antibody (A00084-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for COX2/Cyclooxygenase 2/PTGS2 at approximately 75 kDa. The expected band size for COX2/Cyclooxygenase 2/PTGS2 is at 69 kDa.



IHC analysis of PTGS2 using anti-PTGS2 antibody (A00084-2).

PTGS2 was detected in a paraffin-embedded section of human lung cancer tissue. The tissue section was incubated with rabbit anti-PTGS2 Antibody (A00084-2) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.

Product datasheet
**Anti-COX2/Cyclooxygenase 2/PTGS2
Antibody**

Catalog Number: A00084-2


BOSTER[®]

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com



Flow cytometry analysis of CACO-2 cell(1:100) DyLight488 conjugated goat anti-rabbit IgG(blue) was used as secondary antibody. Isotype control antibody (Green line) was rabbit IgG DyLight488. Unlabelled sample (Red line).