

Basic Information

Product Name	Anti-CD19 Antibody
Gene Name	CD19
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.
Immunogen	E. coli-derived mouse CD19 recombinant protein (Position: R19-M237).
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	95 kDa
Dilution Ratios	Western blot (WB):1:500-2000 ELISA: 1:100-1000


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

B-lymphocyte antigen CD19, also known as CD19 (Cluster of Differentiation 19), is a protein that in humans is encoded by the CD19 gene. It is found on the surface of B-cells, a type of white blood cell. Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. The CD19 gene encodes a cell surface molecule that assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation.

Selected Validation Data



Western blot analysis of CD19 using anti-CD19 antibody (A00154-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Ramos whole cell lysates,

Lane 2: human Daudi whole cell lysates,

Lane 3: rat spleen tissue lysates,

Lane 4: mouse spleen tissue lysates,

Lane 5: mouse RAW264.7 whole cell lysates,

Lane 6: mouse EL4 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-CD19 antigen A03957-Aen affinity purified polyclonal antibody (A00154-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for CD19 at approximately 95 kDa. The expected band size for CD19 is at 61 kDa.