

## Basic Information

Product Name	Anti-GSTP1 Antibody	
Gene Name	GSTP1	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E. coli-derived mouse GST3 / GST pi recombinant protein (Position: P2-Q210).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	23 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 ELISA: 1:100-1000 (Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

## Storage

12 months from date of receipt, -20°C as supplied.


## Background Information

Glutathione S-transferases pi (GSTP1), also known as GST3, is an enzyme that in humans is encoded by the GSTP1 gene. This gene is mapped to 11q13.2. GSTP1 has 7 exons and 6 introns contained within approximately 2.8 kilobases. GSTP1 belongs to Glutathione S-transferases (GSTs) which are a family of enzymes that play an important role in detoxification by catalyzing the conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. GSTP1 presents in all tissues and cells, with the exception of red cells, in which only erythrocyte GST (GSTe) is observed. What's more, GSTP1 is a polymorphic gene encoding active, functionally different GSTP1 variant proteins that are thought to function in xenobiotic metabolism and play a role in susceptibility to cancer, and other diseases.

## Reference

Anti-GSTP1 Antibody被引用在2文献中。

## Selected Validation Data



Western blot analysis of GSTP1 using anti-GSTP1 antibody (A00394). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: mouse lung tissue lysates,

Lane 2: mouse testis tissue lysates,


Lane 3: rat spleen tissue lysates,

Lane 4: human A549 whole cell lysates,

Lane 5: human placenta tissue lysates,

Lane 6: human Hela whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-GSTP1 antigen affinity purified polyclonal antibody (A00394) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for GSTP1 at approximately 23 kDa. The expected band size for GSTP1 is at 23 kDa.



IHC analysis of GSTP1 using anti-GSTP1 antibody (A00394).

GSTP1 was detected in a paraffin-embedded section of mouse kidney tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-GSTP1 Antibody (A00394) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.