antibody and ELISA experts BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Basic Information	
Product Name	Anti-NKG2D/KLRK1 Antibody
Gene Name	KLRK1
Source	Rabbit
Clonality	Polyclonal
lsotype	IgG
Species Reactivity	mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human NKG2D, which shares 76.7% amino acid (aa) sequence identity with both mouse and rat NKG2D.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	25 kDa
Dilution Ratios	Western blot (WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

NKG2D is encoded by KLRK1 gene which is located in the NK-gene complex (NKC) situated on and chromosome 12 in humans. Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calciumdependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Readthrough transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster.

Product datasheet Anti-NKG2D/KLRK1 Antibody Catalog Number: A00661-1

antibody and ELISA experts BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Reference

Anti-NKG2D/KLRK1 Antibody被引用在2文献中。

Selected Validation Data



Western blot analysis of NKG2D/KLRK1 using anti-NKG2D/KLRK1 antibody (A00661-1). The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat lymph tissue lysates, Lane 2: rat spleen tissue lysates,

Lane 3: mouse thymus tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-NKG2D/KLRK1 antigen affinity purified polyclonal antibody (A00661-1) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for NKG2D/KLRK1 at approximately 25 kDa. The expected band size for NKG2D/KLRK1 is at 25 kDa .