

Basic Information

Product Name	Anti-14-3-3 Sigma/SFN Antibody	
Gene Name	SFN	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF, FCM, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E. coli-derived human 14-3-3 sigma recombinant protein (Position: M1-S248).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	28 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-400
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-400
	Flow Cytometry (Fixed):	1:50-200
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000
	(Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Stratifin(SFN), also known as 14-3-3 protein sigma, is strongly induced by gamma irradiation and other DNA-damaging agents. The induction of 14-3-3-sigma is mediated by a p53 -responsive element located 1.8 kb upstream of its transcription start site. The protein, called stratifin, was shown to be diffusely distributed in the cytoplasm and was present in cultured epithelial cells. It was most abundant in tissues enriched in stratified keratinizing epithelium.

Selected Validation Data




Figure 1. Western blot analysis of anti-14-3-3 Sigma/SFN antibody (A01127). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human A431 whole cell lysates,

Lane 2: human HeLa whole cell lysates,

Lane 3: human MCF-7 whole cell lysates,

Lane 4: human Hacat whole cell lysates,

Lane 1: rat skin tissue lysates,

Lane 2: mouse Hepa1-6 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-14-3-3

Sigma/SFN antigen affinity purified polyclonal antibody (A01127)

and probed with a goat anti-rabbit IgG-HRP secondary antibody

(Catalog # BA1054). The signal is developed using ECL Plus

Western Blotting Substrate (Catalog # AR1197). A specific band

was detected for 14-3-3 Sigma/SFN at approximately 28 kDa. The

expected band size for 14-3-3 Sigma/SFN is at 28 kDa.




Figure 2. IHC analysis of 14-3-3 sigma using anti-14-3-3 sigma

antibody (A01127). 14-3-3 sigma was detected in paraffin-

embedded section of human rectal cancer tissues. anti-14-3-3

sigma Antibody (A01127) . Biotinylated goat anti-rabbit IgG was

used as secondary antibody . The tissue section was developed

using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with

DAB as the chromogen.




Figure 4. IF analysis of 14-3-3 sigma using anti- 14-3-3 sigma antibody (A01127). 14-3-3 sigma was detected in paraffin-embedded section of A549 cell. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/mL rabbit anti- cortactin Antibody (A01127) . DyLight488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody Visualize using a fluorescence microscope and filter sets appropriate for the label used.