BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

antibody and FLISA

Basic Information		
Product Name	Anti-FOSB Antibody	
Gene Name	FOSB	
Source	Rabbit	
Clonality	Polyclonal	
lsotype	IgG	
Species Reactivity	human	
Tested Application	IHC, ICC/IF, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
Immunogen	E.coli-derived human FOSB recombinant protein (Position: M1-S332).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Dilution Ratios	Immunohistochemistry (IHC): Immunocytochemistry/Immunofluorescence (ICC/IF): Enzyme linked immunosorbent assay (ELISA): (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or mins is required for the staining of formalin/paraffin sections determined by end user.	

## **Storage**

12 months from date of receipt, -20°C as supplied.


## **Background Information**

FOSB, FBJ murine osteosarcoma viral oncogene homolog B, is a protein that, in humans, is encoded by the FOSB gene. FOSB is a member of Fos gene family which consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. The FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. The FOSB gene is mapped to 19q13.32. Delta FOSB is a truncated splice variant of FOSB. Delta FosB has been implicated in the development of drug addiction and control of the reward system in the brain, and is linked to changes in a number of other gene products such as CREB and sirtuins. Delta FosB also regulates the commitment of mesenchymal precursor cells to the adipocyte or osteoblast lineage.


## **Selected Validation Data**

antibody and ELISA experts BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com



IHC analysis of FOSB using anti-FOSB antibody (A01569-1). FOSB was detected in a paraffin-embedded section of human lung cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-FOSB Antibody (A01569-1) at a dilution of 1:200 and developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.



IF analysis of FOSB using anti-FOSB antibody (A01569-1). FOSB was detected in an immunocytochemical section of SiHa cells. The section was incubated with rabbit anti-FOSB Antibody (A01569-1) at a dilution of 1:100. Dylight594-conjugated Anti-rabbit IgG Secondary Antibody (red)(Catalog#BA1142) was used as secondary antibody. The section was counterstained with DAPI (Catalog # AR1176) (Blue).