Product datasheet

Anti-LAMP2 Antibody

Catalog Number: A01573-3



Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Basic Information	
Product Name	Anti-LAMP2 Antibody
Gene Name	LAMP2
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived human LAMP2 recombinant protein (Position: D128-F410).
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	100-120 kDa
Dilution Ratios	Western blot (WB):1:500-2000 ELISA: 1:100-1000

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Lysosome-associated membrane protein 2 (LAMP2), also known as CD107b (Cluster of Differentiation 107b) and Mac-3, is a human gene. Its protein, LAMP2, is one of the lysosome-associated membrane glycoproteins. The protein encoded by this gene is a member of a family of membrane glycoproteins. This glycoprotein provides selectins with carbohydrate ligands. It may play a role in tumor cell metastasis. It may also function in the protection, maintenance, and adhesion of the lysosome. Alternative splicing of this gene results in multiple transcript variants encoding distinct proteins.

Selected Validation Data


Product datasheet Anti-LAMP2 Antibody Catalog Number: A01573-3



BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com



Western blot analysis of LAMP2 using anti-LAMP2 antibody (A01573-3). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: U-87MG whole cell lysates,

Lane 2: PC-3 whole cell lysates,

Lane 3: 293T whole cell lysates,

Lane 4: Jurkat whole cell lysates,

Lane 5: HepG2 whole cell lysates,

Lane 6: Hela whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-LAMP2 antigen affinity purified polyclonal antibody (A01573-3) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for LAMP2 at approximately 100-120 kDa. The expected band size for LAMP2 is at 46 kDa.