

Basic Information

Product Name	Anti-ATF1 DyLight 488 Conjugated Antibody
Gene Name	ATF1
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human
Tested Application	FCM
Contents	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Immunogen	E.coli-derived human ATF1 recombinant protein (Position: M1-V271). Human ATF1 shares 91% amino acid (aa) sequence identity with mouse ATF1.
Fluorophores	Amax=488nm; Emax=515-545nm
Conjugate	DyLight 488
Concentration	500ug/ml
Purification	Immunogen affinity purified.
Dilution Ratios	Flow cytometry (FCM):1-3 µg/1x10 ⁶ cells

Storage

At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.

Background Information

ATF1, also known as activating transcription factor 1, is a protein that in humans is encoded by the ATF1 gene. It is mapped to 12q13.12. This gene encodes an activating transcription factor, which belongs to the ATF subfamily and bZIP (basic-region leucine zipper) family. It influences cellular physiologic processes by regulating the expression of downstream target genes, which are related to growth, survival, and other cellular activities. This protein is phosphorylated at serine 63 in its kinase-inducible domain by serine/threonine kinases, cAMP-dependent protein kinase A, calmodulin-dependent protein kinase I/II, mitogen- and stress-activated protein kinase and cyclin-dependent kinase 3 (cdk-3). Its phosphorylation enhances its transactivation and transcriptional activities, and enhances cell transformation.

Selected Validation Data

Product datasheet

**Anti-ATF1 DyLight 488 Conjugated
Antibody**

Catalog Number: A01600-Dyl488

BOSTER

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator,
Wuhan East Lake High-tech Development Zone

Web: www.boster.com **Phone:** 027-67845390 **Email:** boster@boster.com

