Product datasheet Anti-BDNF Antibody Catalog Number: BA0565-2

BOSTER®

antibody and ELISA experts
BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Basic Information	
Product Name	Anti-BDNF Antibody
Gene Name	BDNF
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminal of human BDNF, identical to the related rat and mouse sequences.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	28/37 kDa
Dilution Ratios	Western blot (WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Brain derived neurotrophic factor(BDNF) is a member of the neurotrophin family of growth factors that includes NGF, NT3, and NT4. BDNF is a prosurvival factor induced by cortical neurons that is necessary for survival of striatal neurons in the brain. BDNF is expressed within peripheral ganglia and is not restricted to neuronal target fields. BDNF has been purified and shown to reduce the amount of naturally occurring neuronal cell death in portions of the peripheral nervous system.

Reference


Anti-BDNF Antibody被引用在8文献中。



Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Selected Validation Data



Western blot analysis of BDNF using anti-BDNF antibody (BA0565-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Rat brain tissue lysates,

Lane 2: Mouse brain tissue lysates,

Lane 3: MCF-7 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-BDNF antigen affinity purified polyclonal antibody (BA0565-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for BDNF at approximately 28/37 kDa. The expected band size for BDNF is at 28 kDa.