

Basic Information

Product Name	Anti-Caspase 1/CASP1 (p10) Antibody	
Gene Name	CASP1	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Caspase-1(P10).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	45 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-400
	(Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


Background Information

Caspase 1, apoptosis-related cysteine protease, is a cysteine protease that regulates inflammatory processes through its capacity to process and activate the interleukin-1-beta, IL18, and IL33 precursor proteins. Caspase 1 was purified ICE from the cytosol of the THP. human monocytic cell line and found that the active protease was made up of 2 peptides, which they called p20 and p10 based on their apparent molecular masses by SDS-PAGE. It belongs to a family of cysteine proteases known as caspases that always cleave proteins following an aspartic acid residue. The Caspase1 gene consists of 10 exons spanning at least 10.6 kb. The Caspase 1 gene is mapped to 11q23, a site frequently involved in rearrangement in human cancers, including a number of leukemias and lymphomas, by Southern DNA blot analysis of rodent-human hybrids and by in situ hybridization to normal human metaphase chromosomes. Caspase 1 has been shown to induce cell necrosis or pyroptosis and may function in various developmental stages.

Reference

Anti-Caspase 1/CASP1 (p10) Antibody被引用在6文献中。

Selected Validation Data



Western blot analysis of Caspase 1/CASP1 (p10) using anti-Caspase 1/CASP1 (p10) antibody (BA0586). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.


Lane 1: MCF-7 whole cell lysates,

Lane 2: HELA whole cell lysates,

Lane 3: SW620 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-Caspase 1/CASP1 (p10) antigen affinity purified polyclonal antibody (BA0586) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Caspase 1/CASP1 (p10) at approximately 45 kDa. The expected band size for Caspase 1/CASP1 (p10) is at 45 kDa.



IHC analysis of Caspase 1/CASP1 (p10) using anti-Caspase 1/CASP1 (p10) antibody (BA0586).

Caspase 1/CASP1 (p10) was detected in a paraffin-embedded section of human mammary cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-Caspase 1/CASP1 (p10) Antibody (BA0586) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.