

## Basic Information

<b>Product Name</b>	Anti-IkB Alpha/NFKBIA Antibody
<b>Gene Name</b>	NFKBIA
<b>Source</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Species Reactivity</b>	human, mouse, rat, rabbit
<b>Tested Application</b>	IHC
<b>Contents</b>	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.
<b>Immunogen</b>	E.coli-derived human IKB alpha recombinant protein (Position: Q3-Q112). Human IKB alpha shares 87% and 86% amino acid (aa) sequence identity with mouse and rat IKB alpha, respectively.
<b>Concentration</b>	500 ug/ml
<b>Purification</b>	Immunogen affinity purified.
<b>Dilution Ratios</b>	Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


## Background Information

NFKBIA, also called IKBA or MAD-3, is one member of a family of cellular proteins that function to inhibit the NF-κB transcription factor. It is mapped to 14q13.2. NFKBIA inhibits NF-κB by masking the nuclear localization signals(NLS) of NF-κB proteins and keeping them sequestered in an inactive state in the cytoplasm. It moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. The effect of the nonpathogenic bacteria is specific to the SCF complex substrates CTNNB1 and NFKBIA. This may help to explain the beneficial effects of treatment of inflammatory bowel disease with nonpathogenic probiotic enteric organisms. In addition, NFKBIA blocks the ability of NF-κB transcription factors to bind to DNA, which is required for NF-κB's proper functioning.

## Reference

Anti-IkB Alpha/NFKBIA Antibody被引用在5文献中。

## Selected Validation Data



IHC analysis of IkB Alpha/NFKBIA using anti-IkB Alpha/NFKBIA antibody (BA0943-1).

IkB Alpha/NFKBIA was detected in a paraffin-embedded section of human lung cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-IkB Alpha/NFKBIA Antibody (BA0943-1) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.