

Basic Information

Product Name	Anti-GJA5 Antibody	
Gene Name	GJA5	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Connexin 40/GJA5, identical to the related mouse and rat sequences.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	40 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Connexin proteins oligomerize to form intercellular channels, called gap junctions, through which ions and small molecules move between adjacent cells. Connexin 40 (Gap junction alpha-5 protein) is a protein that in humans is encoded by the GJA5 gene. This gene is a member of the connexin gene family. The encoded protein is a component of gap junctions, which are composed of arrays of intercellular channels that provide a route for the diffusion of low molecular weight materials from cell to cell. Mutations in this gene may be associated with atrial fibrillation. Alternatively spliced transcript variants encoding the same isoform have been described.

Reference

Anti-GJA5 Antibody被引用在2文献中。

Selected Validation Data




Western blot analysis of GJA5 using anti-GJA5 antibody (BA1593-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Mouse Heart tissue lysates,

Lane 2: Mouse Heart tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-GJA5 antigen affinity purified polyclonal antibody (BA1593-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for GJA5 at approximately 40 kDa. The expected band size for GJA5 is at 40 kDa.



IHC analysis of GJA5 using anti-GJA5 antibody (BA1593-2).

GJA5 was detected in a paraffin-embedded section of rat cardiac muscle tissue. The tissue section was incubated with rabbit anti-GJA5 Antibody (BA1593-2) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.