

## Basic Information

<b>Product Name</b>	Anti-NPHS2 Antibody
<b>Gene Name</b>	NPHS2
<b>Source</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Species Reactivity</b>	human, mouse, rat
<b>Tested Application</b>	WB
<b>Contents</b>	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.
<b>Immunogen</b>	A synthetic peptide corresponding to a sequence in the middle region of human NPHS2, different from the mouse sequence by one amino acid.
<b>Concentration</b>	500 ug/ml
<b>Purification</b>	Immunogen affinity purified.
<b>Observed MW</b>	42 kDa
<b>Dilution Ratios</b>	Western blot (WB):1:500-2000

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

## Background Information

Podocin(PDCN) is a protein which lines the podocytes and assists in maintaining the barrier at the glomerular basement membrane. NPHS2 is a causative gene for Familial idiopathic nephrotic syndromes, which represents a heterogeneous group of kidney disorders, and include autosomal recessive steroid-resistant nephrotic syndrome, which is characterized by early childhood onset of proteinuria, rapid progression to end-stage renal disease and focal segmental glomerulosclerosis. By positional cloning, NPHS2 was mapped to 1q25-31. It is almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli, and encodes a new integral membrane protein, podocin, belonging to the stomatin protein family. Boute et al.(2000) found ten different NPHS2 mutations, comprising nonsense, frameshift and missense mutations, to segregate with the disease, demonstrating a crucial role for podocin in the function of the glomerular filtration barrier.

## Reference

Anti-NPHS2 Antibody被引用在1文献中。

## Selected Validation Data



Figure 1. Western blot analysis of NPHS2 using anti-NPHS2 antibody (BA1688-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat kidney tissue lysates,

Lane 2: human Hela whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-NPHS2 antigen affinity purified polyclonal antibody (BA1688-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for NPHS2 at approximately 42 kDa. The expected band size for NPHS2 is at 42 kDa.