

Basic Information

Product Name	Anti-NFkB/NFKB2 p100/p52 Antibody	
Gene Name	NFKB2	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	mouse, rat	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of mouse NFkB/NFKB2 p100/p52, identical to the related rat sequence.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	52 kDa(activeform)/120 kDa(precursor)	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied.

Background Information


NFKB2(Nuclear Factor Kappa-B, Subunit 2), also known as NFKB or p52/p100 SUBUNIT, is a protein that in humans is encoded by the NFKB2 gene. Liptay et al.(1992) mapped the gene for what they called the p49/p100 subunit of NFKB(NFKB2) to chromosome 10 by Southern blot analysis of panels of human/Chinese hamster cell hybrids. By fluorescence in situ hybridization(FISH), they confirmed the localization and mapped the gene with greater resolution to 10q24. NFKB2 appears to be the same as LYT10. Claudio et al.(2002) showed that bone marrow(BM) cells from Nfkb2-deficient mice, but not Nfkb1-deficient mice, failed to increase relative and total IgD-positive transitional-1(T1) stage B cells in response to Baff. In vivo, however, Nfkb2-deficient mice did generate mature B cells, but at reduced numbers. Mice of the aly/aly strain, which are naturally deficient in Nik, and mice of the A/WySNJ strain, which have a mutation in Baffr, also failed to produce T1 B cells in response to Baff. Baff stimulation enhanced expression of Bcl2 in T1 B cells, thereby promoting B-cell survival, and caused the processing of the p100 form of Nfkb2 to p52, which again required Baffr and Nik, but not Nemo(IKKG). Immunoblot analysis showed that BM cells contained


primarily p100.

Reference

Anti-NFkB/NFKB2 p100/p52 Antibody被引用在3文献中。

Selected Validation Data





IHC analysis of NFkB/NFKB2 p100/p52 using anti-NFkB/NFKB2 p100/p52 antibody (BA1873).

NFkB/NFKB2 p100/p52 was detected in a paraffin-embedded section of rat intestine tissue. The tissue section was incubated with rabbit anti-NFkB/NFKB2 p100/p52 Antibody (BA1873) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.