

## Basic Information

<b>Product Name</b>	Anti-PAPPA Antibody
<b>Gene Name</b>	PAPPA
<b>Source</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Species Reactivity</b>	human, mouse, rat, rabbit
<b>Tested Application</b>	IHC
<b>Contents</b>	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.
<b>Immunogen</b>	E.coli-derived human PAPP A recombinant protein (Position: R95-Q388). Human PAPP A shares 88% amino acid (aa) sequence identity with mouse PAPP A.
<b>Concentration</b>	500 ug/ml
<b>Purification</b>	Immunogen affinity purified.
<b>Dilution Ratios</b>	Immunohistochemistry in paraffin section IHC-(P): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


## Background Information

Pappalysin-1, also known as DIPLA1, is a protein that in humans is encoded by the PAPPA gene. It is mapped to 9q33.1. PAPPA is found in the ovarian follicles, follicular fluid, luteal cells, and fallopian tubes of nonpregnant women and in the seminal vesicles and seminal fluid of males. This gene encodes a secreted metalloproteinase which cleaves insulin-like growth factor binding proteins (IGFBPs). It is thought to be involved in local proliferative processes such as wound healing and bone remodeling. Low plasma level of this protein has been suggested as a biochemical marker for pregnancies with aneuploid fetuses. It has been found that circulating PAPPA is a disulfide-bridged complex with proMBP in which the subunits of the constituents are present in a 1:1 molar ratio.

## Reference

Anti-PAPPA Antibody被引用在1文献中。

## Selected Validation Data



IHC analysis of PAPPA using anti-PAPPA antibody (BA2327-1). PAPPA was detected in a paraffin-embedded section of human placenta tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-PAPPA Antibody (BA2327-1) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.