

Basic Information

Product Name	Anti-Survivin/BIRC5 Antibody (Clone#ABB-2)	
Gene Name	BIRC5	
Source	Rabbit	
Clonality	Monoclonal	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, IHC, ICC/IF, IP, FCM	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
Immunogen	A synthesized peptide derived from human Survivin	
Concentration	500 ug/ml	
Purification	Affinity-chromatography	
Observed MW	16 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-200
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-200
	ImmunoPrecipitation (IP):	1:20
	Flow Cytometry (FCM):	1:20

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Survivin, also called baculoviral inhibitor of apoptosis repeat-containing 5 or BIRC5, is a protein that, in humans, is encoded by the BIRC5 gene. Survivin is a member of the inhibitor of apoptosis(IAP) family. The survivin gene contains 4 exons. The survivin gene is mapped to chromosome 17q25 by pulsed field gel electrophoresis and single- and 2-color FISH. The survivin protein functions to inhibit caspase activation, thereby leading to negative regulation of apoptosis or programmed cell death. The survivin protein is expressed highly in most human tumours and fetal tissue, but is completely absent in terminally differentiated cells.

Reference

Anti-Survivin/BIRC5 Antibody (Clone#ABB-2)被引用在2文献中。

Selected Validation Data



Western blot analysis of anti-Survivin/BIRC5 antibody (BM3975). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human K562 whole cell lysates,


Lane 3: human Raji whole cell lysates,

Lane 4: human PC-3 whole cell lysates,

Lane 5: human U2OS whole cell lysates,


Lane 6: human 293T whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-Survivin/BIRC5 antigen affinity purified monoclonal antibody (BM3975) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Survivin/BIRC5 at approximately 16 kDa. The expected band size for Survivin/BIRC5 is at 16 kDa.



Immunohistochemical analysis of paraffin-embedded human gastric carcinoma, using Survivin Antibody.

Hela



Immunofluorescent analysis using the Antibody.