

Basic Information

Product Name	Anti-P27/KIP1/CDKN1B Antibody (Clone#CGA-3)	
Gene Name	CDKN1B	
Source	Rabbit	
Clonality	Monoclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF, IP, FCM	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
Immunogen	A synthesized peptide derived from human p27 KIP 1	
Concentration	500 ug/ml	
Purification	Affinity-chromatography	
Observed MW	25 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-200 Immunocytochemistry/Immunofluorescence (ICC/IF): 1:50-200 ImmunoPrecipitation (IP): 1:50 Flow Cytometry (FCM): 1:50	

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

P27(kip1), also known as cyclin dependent kinase inhibitor 1B(CDKN1B), is a major target of AFX-like forkhead proteins. CDKN1B(p27) belongs to the Cip/Kip family and functions as an important cell cycle gatekeeper. Phosphorylation leads to the ubiquitination and degradation of CDKN1B. P27(kip1) mapped to 12p13. An increase in p27 causes proliferating cells to exit from the cell cycle, and a decrease in p27 is necessary for quiescent cells to resume division. Abnormally low amounts of p27 are associated with pathological states of excessive cell proliferation, especially cancers. Overexpression of p27Kip1 lengthens the G1 phase in a mouse model that targets inducible gene expression to central nervous system progenitor cells.

Reference

Anti-P27/KIP1/CDKN1B Antibody (Clone#CGA-3)被引用在3文献中。

Selected Validation Data



Western blot analysis of anti-P27/KIP1/CDKN1B antibody (BM4229). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human MCF-7 whole cell lysates,

Lane 2: human Jurkat whole cell lysates,


Lane 3: human Caco-2 whole cell lysates,

Lane 4: human HepG2 whole cell lysates,

Lane 5: rat brain tissue lysates,


Lane 6: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-P27/KIP1/CDKN1B antigen affinity purified monoclonal antibody (BM4229) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for P27/KIP1/CDKN1B at approximately 25 kDa. The expected band size for P27/KIP1/CDKN1B is at 22 kDa.



IHC analysis of P27/KIP1/CDKN1B using anti-P27/KIP1/CDKN1B antibody (BM4229).

P27/KIP1/CDKN1B was detected in a paraffin-embedded section of human breast cancer tissue. The tissue section was incubated with rabbit anti-P27/KIP1/CDKN1B Antibody (BM4229) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.



Immunofluorescent analysis of Hela cells, using p27 KIP 1 Antibody .