

Basic Information

Product Name	Anti-PARK2/Parkin/PRKN Antibody (Clone#AOCD-16)	
Gene Name	PRKN	
Source	Rabbit	
Clonality	Monoclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF, IP, FCM	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
Immunogen	A synthesized peptide derived from human Parkin	
Concentration	500 ug/ml	
Purification	Affinity-chromatography	
Observed MW	52,66 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-200
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-200
	ImmunoPrecipitation (IP):	1:20
	Flow Cytometry (FCM):	1:20

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Parkin is a RING domain-containing E3 ubiquitin ligase involved in proteasome-dependent degradation of proteins. It is mapped to 6q26. This gene is important for mitochondrial quality control by lysosome-dependent degradation of damaged mitochondria through autophagy, or mitophagy. Parkin is expressed in neuronal processes and cell bodies of neurons, but not glial cells, in the midbrain, basal ganglia, cerebral cortex, and cerebellum. Parkin assimilated with actin filaments, suggesting that it is a cytoskeletal-associated protein. Parkin is identified as a transcriptional repressor of p53 independent of its ubiquitin ligase function. It also has been found that parkin was associated physically with mitochondrial DNA (mtDNA) in proliferating as well as in differentiated SH-SY5Y neuroblastoma cells.

Reference

Anti-PARK2/Parkin/PRKN Antibody (Clone#AOCD-16)被引用在7文献中。

Selected Validation Data

Western blot analysis of Parkin expression in Jurkat cell lysate.

