

## Basic Information

<b>Product Name</b>	Anti-CD68 Antibody (Clone#SP130)
<b>Gene Name</b>	CD68
<b>Source</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Species Reactivity</b>	human
<b>Tested Application</b>	IF,IHC
<b>Contents</b>	200 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol.
<b>Immunogen</b>	Subcellular fraction of human alveolar macrophages
<b>Concentration</b>	200 ug/ml
<b>Dilution Ratios</b>	Immunohistochemistry (IHC):1:50-100 Immunofluorescence (IF): 1:50-100

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

## Background Information

CD68, cluster of differentiation, is a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. CD68 is a member of a family of hematopoietic mucin-like molecules that includes leukosialin/CD43 and stem cell antigen CD34. The CD68 gene is mapped to 17p13.1. Immunohistochemistry can be used to identify the presence of CD68, which is found in the cytoplasmic granules of a range of different blood cells. It is particularly useful as a marker for the various cells of the macrophage lineage, including monocytes, histiocytes, giant cells, Kupffer cells, and osteoclasts. This allows it to be used to distinguish diseases of otherwise similar appearance, such as the monocyte/macrophage and lymphoid forms of leukaemia (the latter being CD68 negative). Its presence in macrophages also makes it useful in diagnosing conditions related to proliferation or abnormality of these cells, such as malignant histiocytosis, histiocytic lymphoma, and Gaucher's disease.

## Reference

Anti-CD68 Antibody (Clone#SP130)被引用在9文献中。

Product datasheet

## Anti-CD68 Antibody (Clone#SP130)

Catalog Number: **BM5785**

# BOSTER

antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator,  
Wuhan East Lake High-tech Development Zone

**Web:** [www.boster.com](http://www.boster.com) **Phone:** 027-67845390 **Email:** [boster@boster.com](mailto:boster@boster.com)

## Selected Validation Data




Figure 1. IF analysis using anti- CD68 antibody (BM5785). detected in paraffin-embedded section of human tonsil tissues. The tissue section were stained using the cy3-conjugated Anti-mouse IgG Secondary Antibody (red)(Catalog # BA1031).