

Basic Information

Product Name	Anti-KGA/GAC/GLS Antibody (Clone#9G6)	
Gene Name	GLS	
Source	Mouse	
Clonality	Monoclonal	
Isotype	IgG2b	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E. coli-derived human Glutaminase/GLS recombinant protein (Position: K396-N654). Human GLS shares 99.6% amino acid (aa) sequence identity with both mouse and rat GLS.	
Concentration	500 ug/ml	
Purification	protein G purified.	
Observed MW	65-73 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 Immunocytochemistry/Immunofluorescence (ICC/IF): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

This gene encodes the K-type mitochondrial glutaminase. The encoded protein is an phosphate-activated amidohydrolase that catalyzes the hydrolysis of glutamine to glutamate and ammonia. This protein is primarily expressed in the brain and kidney plays an essential role in generating energy for metabolism, synthesizing the brain neurotransmitter glutamate and maintaining acid-base balance in the kidney. Alternate splicing results in multiple transcript variants.

Selected Validation Data

Product datasheet
Anti-KGA/GAC/GLS Antibody
(Clone#9G6)

Catalog Number: M01272-2




antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com




Western blot analysis of KGA/GAC/GLS using anti-KGA/GAC/GLS antibody (M01272-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: HepG2 whole cell lysates,


Lane 2: HEL whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with mouse anti-KGA/GAC/GLS antigen affinity purified monoclonal antibody (M01272-2) at a dilution of 1:1000 and probed with a goat anti-mouse IgG-HRP secondary antibody (Catalog # BA1050). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for KGA/GAC/GLS at approximately 65-73 kDa. The expected band size for KGA/GAC/GLS is at 73 kDa.



IHC analysis of KGA/GAC/GLS using anti-KGA/GAC/GLS antibody (M01272-2).

KGA/GAC/GLS was detected in a paraffin-embedded section of human Gall bladder adenosquamous carcinoma tissue. Biotinylated goat anti-mouse IgG was used as secondary antibody. The tissue section was incubated with mouse anti-KGA/GAC/GLS Antibody (M01272-2) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB (Catalog # AR1027) as the chromogen.



IF analysis of KGA/GAC/GLS using anti-KGA/GAC/GLS antibody (M01272-2). KGA/GAC/GLS was detected in an immunocytochemical section of SiHa cells. The section was incubated with mouse anti-KGA/GAC/GLS Antibody (M01272-2) at a dilution of 1:100. Dylight488-conjugated Anti-mouse IgG Secondary Antibody (green)(Catalog#BA1126) was used as secondary antibody. The section was counterstained with DAPI (Catalog # AR1176) (Blue).