

## Basic Information

<b>Product Name</b>	Anti-FGF2 Antibody (Clone#OTI3D9)	
<b>Gene Name</b>	FGF2	
<b>Source</b>	Mouse	
<b>Clonality</b>	Monoclonal	
<b>Isotype</b>	IgG2b	
<b>Species Reactivity</b>	human, mouse, rat	
<b>Tested Application</b>	WB, IHC, ICC/IF	
<b>Contents</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
<b>Immunogen</b>	Human recombinant protein fragment corresponding to amino acids 10-155 of human bFGF (NP_001997) produced in E.coli.	
<b>Concentration</b>	500 ug/ml	
<b>Purification</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
<b>Observed MW</b>	31 kDa	
<b>Dilution Ratios</b>	Western blot (WB): 1:1000 Immunohistochemistry (IHC): 1:50 Immunocytochemistry/Immunofluorescence (ICC/IF):1:100	

## Storage

Stable for 12 months from date of receipt. Store at -20°C as received.


## Background Information

FGF2 has been implicated in a multitude of physiologic and pathologic processes, including limb development, angiogenesis, wound healing, and tumor growth. Human FGF2 shares 96% and 97% amino acid sequence homology with mouse and rat respectively. FGF2 belongs to the fibroblast growth factor (FGF) family. Fibroblast growth factors (FGFs) exhibit widespread mitogenic and neurotrophic activities. Nine members of the family are currently known, and FGF-1 and FGF-2 are present in relatively high levels in CNS. FGF-2 is expressed by at low levels in many tissues and cell types and reaches high concentrations in brain and pituitary.


## Reference

Anti-FGF2 Antibody (Clone#OTI3D9)被引用在1文献中。


## Selected Validation Data



Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-BFGF monoclonal antibody at 1:200 dilution.



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-BFGF mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, MA00121)



Anti-BFGF mouse monoclonal antibody immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BFGF .