

Basic Information

Product Name	Anti-NR4A1 Antibody
Gene Name	NR4A1
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human NUR77 different from the related mouse sequence by one amino acid, and from the related rat sequence by two amino acids.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	67 kDa
Dilution Ratios	Western blot (WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

NR4A1 (NUCLEAR RECEPTOR SUBFAMILY 4, GROUP A, MEMBER 1), also called NAK1, GFRP1, TR3, NUR77 or NGFIB, is a protein that in humans is encoded by the NR4A1 gene, which is also a member of the Nur nuclear receptor family of intracellular transcription factors. The NR4A1 gene is mapped on 12q13.13. NR4A1 is involved in cell cycle mediation, inflammation and apoptosis. It plays a key role in mediating inflammatory responses in macrophages. In addition, subcellular localization of the NR4A1 protein appears to play a key role in the survival and death of cells. Nr4a1 is overexpressed in Wnt1 -transformed mouse mammary cells. Nr4a1 is also induced by lithium, a Wnt1 mimic, and the Nr4a1 promoter is activated by lithium and beta-catenin, a Wnt1 downstream effector. In contrast, human NR4A1 is not upregulated by beta-catenin, indicating that this gene is regulated differently in human and mouse cells. Adenoviral expression of Nr4a1 induces genes involved in gluconeogenesis, stimulates glucose production both in vitro and in vivo, and raises blood glucose levels.

Reference

Anti-NR4A1 Antibody被引用在1文献中。

Selected Validation Data



Western blot analysis of NR4A1 using anti-NR4A1 antibody (PB0818). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Rat Brain tissue lysates,

Lane 2: Mouse Brain tissue lysates,

Lane 3: U87 whole cell lysates,

Lane 4: HELA whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-NR4A1 antigen affinity purified polyclonal antibody (PB0818) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for NR4A1 at approximately 67 kDa. The expected band size for NR4A1 is at 64 kDa.