

Basic Information

Product Name	Anti-MIF Antibody	
Gene Name	MIF	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, IHC, ICC/IF, FCM	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E. coli-derived human MIF recombinant protein(Position: M1-A115).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	12 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 Immunocytochemistry/Immunofluorescence (ICC/IF): 1:50-400 Flow Cytometry (Fixed): 1:50-200 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


Background Information

Macrophage migration inhibitory factor, MIF, is a cytokine released by T-lymphocytes, macrophages, and the pituitary gland that serves to integrate peripheral and central inflammatory responses. MIF gene has 3 exons separated by introns of only 189 and 95 bp, and covers less than 1kb. Localization of the human gene for macrophage migration inhibitory factor(MIF) to chromosome 22q11.2. MIF plays a critical role in inflammatory diseases and atherogenesis.

Reference

Anti-MIF Antibody被引用在1文献中。

Selected Validation Data



Western blot analysis of anti-MIF antibody (PB9274). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human K562 whole cell lysates,

Lane 3: human Hela whole cell lysates,


Lane 4: human HepG2 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-MIF antigen affinity purified polyclonal antibody (PB9274) and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054).

The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for MIF at

approximately 12 kDa. The expected band size for MIF is at 12 kDa.




IHC analysis of MIF using anti-MIF antibody (PB9274).

MIF was detected in a paraffin-embedded section of human liver

cancer tissue. The tissue section was developed using HRP


Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002)

with DAB (Catalog # AR1027) as the chromogen.



IF analysis of MIF using anti-MIF antibody (PB9274).

MIF was detected in an immunocytochemical section of Hela cells. DyLight®488 Conjugated Goat Anti-Rabbit IgG (Green) (Catalog # BA1127) was used as secondary antibody. The section was counterstained with DAPI (Catalog # AR1176) (Blue).



Flow Cytometry analysis of Jurkat cells using anti-MIF antibody (PB9274).

Overlay histogram showing Jurkat cells stained with PB9274 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MIF Antibody (PB9274, 1:100).

DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 1:100) was used as secondary antibody. Isotype control antibody (Green line) was rabbit IgG (Catalog # BA1045) (1:100) used under the same conditions. Unlabelled sample (Red line) was also used as a control.