

## Basic Information

Product Name	Anti-CYP19A1 Antibody
Gene Name	CYP19A1
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived mouse Aromatase/Cyp19a1 recombinant protein (Position: N75-Q428).
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	58 kDa
Dilution Ratios	Western blot (WB):1:500-2000 ELISA: 1:100-1000

## Storage

12 months from date of receipt, -20°C as supplied.


## Background Information

CYP19A1, also called Aromatase, is an enzyme responsible for a key step in the biosynthesis of estrogens. It is a member of the cytochrome P450 superfamily, which are monooxygenases that catalyze many reactions involved in steroidogenesis. In particular, aromatase is responsible for the aromatization of androgens into estrogens. The CYP19 gene spans at least 70 kb of genomic DNA and contains 10 exons. By in situ hybridization, the ARO gene is mapped to 15q21.1. The aromatase enzyme can be found in many tissues including gonads, brain, adipose tissue, placenta, blood vessels, skin, bone, and endometrium, as well as in tissue of endometriosis, uterine fibroids, breast cancer, and endometrial cancer. It is an important factor in sexual development. Some bodybuilders taking steroids also take antiaromatase supplements to prevent excess testosterone conversion into estrogens, which can cause gynecomastia.

## Reference

Anti-CYP19A1 Antibody 被引用在4文献中。

## Selected Validation Data



Western blot analysis of CYP19A1 using anti-CYP19A1 antibody

(A00071-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human placenta tissue lysates,

Lane 2: HepG2 whole cell lysates,

Lane 3: HCCT tissue lysates,

Lane 4: HCCP tissue lysates,

Lane 5: U-87MG whole cell lysates,

Lane 6: MCF-7 whole cell lysates,

Lane 7: SK-OV-3 whole cell lysates,

Lane 8: SIHA whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-CYP19A1 antigen affinity purified polyclonal antibody (A00071-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for CYP19A1 at approximately 58 kDa. The expected band size for CYP19A1 is at 58 kDa.