

Basic Information

Product Name	Anti-Bcl-XL/BCL2L1 Antibody
Gene Name	BCL2L1
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, FCM
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived human Bcl-XL recombinant protein (Position: M1-T219). Human Bcl-XL shares 97.9% amino acid (aa) sequence identity with both mouse and rat Bcl-XL.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	26 kDa、19 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Flow Cytometry (Fixed):1:50-200

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


Background Information

Bcl-2-like protein 1, also known as Bcl-XL, is a protein that in humans is encoded by the BCL2L1 gene. The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform (Bcl-xL) acts as an apoptotic inhibitor and the shorter form (Bcl-xS) acts as an apoptotic activator.

Reference

Anti-Bcl-XL/BCL2L1 Antibody被引用在4文献中。

Selected Validation Data



Western blot analysis of Bcl-XL/BCL2L1 using anti-Bcl-XL/BCL2L1 antibody (A00181). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human Hela whole cell lysates,

Lane 3: human A549 whole cell lysates,

Lane 4: human Raji whole cell lysates,


Lane 5: rat brain tissue lysates,

Lane 6: rat C6 whole cell lysates,

Lane 7: mouse brain tissue lysates,

Lane 8: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-Bcl-XL/BCL2L1 antigen affinity purified polyclonal antibody (A00181) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Bcl-XL/BCL2L1 at approximately 26-30 kDa. The expected band size for Bcl-XL/BCL2L1 is at 26 kDa.



Flow Cytometry analysis of Jurkat cells using anti-Bcl-XL/BCL2L1 antibody (A00181).

Overlay histogram showing Jurkat cells stained with A00181 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Bcl-XL/BCL2L1 Antibody (A00181) at 1:100 dilution for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Product datasheet

Anti-Bcl-XL/BCL2L1 Antibody

Catalog Number: **A00181**



antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com