

## Basic Information

Product Name	Anti-BAFF/TNFSF13B Antibody	
Gene Name	TNFSF13B	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	FCM, WB, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E.coli-derived human BAFF/TNFSF13B recombinant protein (Position: A134-T277).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	31 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Flow Cytometry (Fixed):	1:50-200
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000

## Storage

12 months from date of receipt, -20°C as supplied.


## Background Information

BAFF was regularly detected by enzyme-linked immunosorbent assay in brain tissue lysates and in normal spinal fluid, and in astrocytes by double fluorescence microscopy. BAFF was localized in astrocytes close to BAFF-R-expressing immune cells. BAFF receptors were strongly expressed in situ in primary central nervous system (CNS) lymphomas.<sup>1</sup> The TNF superfamily member B cell-activating factor (BAFF) plays an important role in humoral immunity and in autoimmune diseases, including RA. Local BAFF gene targeting inhibited proinflammatory cytokine expression, suppressed generation of plasma cells and Th17 cells, and markedly ameliorated joint pathology. The B cell activating factor BAFF (BlyS/TALL-1/zTNF4) is a tumor necrosis factor (TNF)-related ligand that promotes B cell survival and binds to three receptors (BCMA, TACI, and the recently described BAFF-R). Human BAFF was mapped to chromosome 13q32-34. The standard used in this kit is recombinant soluble human BAFF (A134-L295) with the molecular mass of 19.6KDa.

## Reference


Anti-BAFF/TNFSF13B Antibody被引用在1文献中。

## Selected Validation Data



Flow Cytometry analysis of HL-60 cells using anti-BAFF/TNFSF13B antibody (A01257).

Overlay histogram showing HL-60 cells stained with A01257 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-BAFF/TNFSF13B Antibody (A01257) at 1:100 dilution for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.



Western blot analysis of BAFF/TNFSF13B using anti-BAFF/TNFSF13B antibody (A01257). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human U-937 whole cell lysates,

Lane 2: human Caco-2 whole cell lysates,

Lane 3: human PANC-1 whole cell lysates,

Lane 4: human CCRF-CEM whole cell lysates,

Lane 5: human MDA-MB-231 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-BAFF/TNFSF13B antigen affinity purified polyclonal antibody (A01257) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for BAFF/TNFSF13B at approximately 31 kDa. The expected band size for BAFF/TNFSF13B is at 31 kDa.