

Basic Information

Product Name	Anti-IDE DyLight 488 Conjugated Antibody
Gene Name	IDE
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human
Tested Application	FCM
Contents	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Immunogen	E. coli-derived human IDE recombinant protein (Position: F485-K756).
Fluorophores	Amax=488nm; Emax=515-545nm
Conjugate	DyLight 488
Concentration	500ug/ml
Purification	Immunogen affinity purified.
Dilution Ratios	Flow cytometry (FCM):1-3 µg/1x10 ⁶ cells

Storage

At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.

Background Information

Insulin-degrading enzyme, also known as IDE, is an enzyme. This gene encodes a zinc metallopeptidase that degrades intracellular insulin, and thereby terminates insulin's activity, as well as participating in intercellular peptide signalling by degrading diverse peptides such as glucagon, amylin, bradykinin, and kallidin. The preferential affinity of this enzyme for insulin results in insulin-mediated inhibition of the degradation of other peptides such as beta-amyloid. Deficiencies in this protein's function are associated with Alzheimer's disease and type 2 diabetes mellitus but mutations in this gene have not been shown to be causative for these diseases. This protein localizes primarily to the cytoplasm but in some cell types localizes to the extracellular space, cell membrane, peroxisome, and mitochondrion. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Selected Validation Data

Product datasheet

Anti-IDE DyLight 488 Conjugated Antibody

Catalog Number: A01358-Dyl488



antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com

暂无图片