Product datasheet Anti-LC3A/MAP1LC3A Antibody Catalog Number: A01543-2

BOSTER®
antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY
Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,

East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Basic Information	
Product Name	Anti-LC3A/MAP1LC3A Antibody
Gene Name	MAP1LC3A
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human MAP1LC3A, identical to the related mouse and rat sequences.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	18 kDa

Western blot (WB):1:500-2000

Storage

Dilution Ratios

12 months from date of receipt, -20°C as supplied.

Background Information

Microtubule-associated proteins 1A/1B light chain 3A is a protein that in humans is encoded by the MAP1LC3A gene. MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. The protein encoded by this gene is one of the light chain subunits and can associate with either MAP1A or MAP1B. Two transcript variants encoding different isoforms have been found for this gene. The expression of variant 1 is suppressed in many tumor cell lines, suggesting that may be involved in carcinogenesis.

Reference

Anti-LC3A/MAP1LC3A Antibody被引用在2文献中。

Product datasheet Anti-LC3A/MAP1LC3A Antibody


BOSTER BIOLOGICAL TECHNOLOGY Catalog Number: A01543-2



Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Selected Validation Data



Western blot analysis of LC3A/MAP1LC3A using anti-LC3A/MAP1LC3A antibody (A01543-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,

Lane 2: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-LC3A/MAP1LC3A antigen affinity purified polyclonal antibody (A01543-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for LC3A/MAP1LC3A at approximately 18 kDa. The expected band size for LC3A/MAP1LC3A is at 14 kDa.