

Basic Information

Product Name	Anti-JUNB Antibody	
Gene Name	JUNB	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, IHC, FCM, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
Immunogen	E.coli-derived human JUNB recombinant protein (Position: T3-F347).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	42-45 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-400
	Flow Cytometry (Fixed):	1:50-200
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Transcription factor jun-B (JUNB) is a protein that in humans is encoded by the JUNB gene. It is mapped to 19p13.2. JUNB is a transcription factor involved in regulating gene activity following the primary growth factor response. It binds to the DNA sequence 5'-TGA[CG]TCA-3', and a large fraction (over 50%) of the JUNB locus is contained in these flanking evolutionarily conserved sequences (FECS), which may be required for effecting the proper transcriptional regulation of this gene. What's more, the expression of JUNB gene might be involved in terminal granulocyte differentiation or in regulating granulocyte functionality.

Selected Validation Data



Western blot analysis of JUNB using anti-JUNB antibody (A01825-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human hela whole cell lysates,

Lane 2: human K562 whole cell lysates,

Lane 3: human CACO-2 whole cell lysates,


Lane 4: human A549 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-JUNB antigen

affinity purified polyclonal antibody (A01825-2) at a dilution of


1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for JUNB at approximately 42-45 kDa. The expected band size for JUNB is at 36 kDa.



IHC analysis of JUNB using anti-JUNB antibody (A01825-2).

JUNB was detected in a paraffin-embedded section of human

mammary cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-JUNB Antibody (A01825-2) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.



Flow Cytometry analysis of U87 cells using anti-JUNB antibody (A01825-2).

Overlay histogram showing U87 cells stained with A01825-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-JUNB Antibody (A01825-2) at 1:100

dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.