BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

antibody and FLISA

Basic Information	
Product Name	Anti-ALOX12 Antibody
Gene Name	ALOX12
Source	Rabbit
Clonality	Polyclonal
lsotype	IgG
Species Reactivity	mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human 12 Lipoxygenase, different from the related mouse sequence by six amino acids, and from the related rat sequence by seven amino acids.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	75 kDa
Dilution Ratios	Western blot (WB):1:500-2000

## **Storage**

12 months from date of receipt, -20°C as supplied.

## **Background Information**


ALOX12, Arachidonate 12-lipoxygenase, is an enzyme that in humans is encoded by the ALOX12 gene. By fluorescence in situ hybridization, the ALOX12 gene is located in band 17p13.1. The gene consists of 14 exons with 13 introns and spans approximately 15 kb of DNA Arachidonate 12-lipoxygenase introduces a molecular oxygen into the C-12 position of arachidonic acid to produce 12(S)-hydroperoxy-5,8,10,14-eicosatetraenoic acid. The major pathway of arachidonic acid metabolism in human platelets proceeds via a 12-lipoxygenase enzyme. Expression of the LOG12 gene was detected in human erythroleukemia cells, platelets, and human umbilical vein endothelial cells by reverse transcription-PCR analysis.

## **Selected Validation Data**

## Product datasheet Anti-ALOX12 Antibody Catalog Number: A02275-1

antibody and ELISA experts BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com



Western blot analysis of ALOX12 using anti-ALOX12 antibody (A02275-1). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat spleen tissue lysates,

Lane 2: mouse spleen tissue lysates,

Lane 3: COLO320 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-ALOX12 antigen affinity purified polyclonal antibody (A02275-1) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for ALOX12 at approximately 75 kDa. The expected band size for ALOX12 is at 76 kDa.