

Basic Information

Product Name	Anti-APEX2 Antibody
Gene Name	APEX2
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, FCM
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived human APEX2 recombinant protein (Position: L102-A210). Human APEX2 shares 91.7% amino acid (aa) sequence identity with mouse APEX2.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	57 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Flow Cytometry (Fixed):1:50-200

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

APEX2, also called apurinic/apyrimidinic endonuclease like-2, is a member of the apurinic/apyrimidinic (AP) family of endonucleases that initiate the repair of AP sites formed by spontaneous hydrolysis of the N-glycosylic bond, mutagen-induced base release, or damaged-base excision by a DNA repair glycosylase. RT-PCR detected APEX2 expression in HeLa cells, Jurkat cells, and human kidney, brain and fetal brain tissue. The APEX2 gene is mapped to chromosome Xp11.21. APEX2 participates in both nuclear and mitochondrial base excision repair (BER) and it can play a role in processing 3-prime-damaged termini or 3-prime-mismatched nucleotides. Additionally, APEX2 displayed weaker AP site-specific and 3-prime nuclease activities compared to APEX1.


Reference

Anti-APEX2 Antibody被引用在1文献中。

Selected Validation Data



After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-APEX2 antigen affinity purified polyclonal antibody (A07203) and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for APEX2 at approximately 57 kDa. The expected band size for APEX2 is at 57 kDa.



Flow Cytometry analysis of A549 cells using anti-APEX2 antibody (A07203).

Overlay histogram showing A549 cells stained with A07203 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-APEX2 Antibody (A07203, 1:100). DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 1:100) was used as secondary antibody. Isotype control antibody (Green line) was rabbit IgG (Catalog # BA1045) (1:100) used under the same conditions. Unlabelled sample (Red line) was also used as a control.