

Basic Information

Product Name	Anti-GRK2 Antibody	
Gene Name	GRK2	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat, monkey	
Tested Application	WB, ICC/IF, FCM	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human GRK2, identical to the related mouse and rat sequences.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	80 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunocytochemistry/Immunofluorescence (ICC/IF): 1:50-400 Flow Cytometry (Fixed): 1:50-200	


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Beta adrenergic receptor kinase (also referred to as β ARK or BARK) is a serine/threonine intracellular kinase. The product of this gene phosphorylates the beta-2-adrenergic receptor and appears to mediate agonist-specific desensitization observed at high agonist concentrations. This protein is an ubiquitous cytosolic enzyme that specifically phosphorylates the activated form of the beta-adrenergic and related G-protein-coupled receptors. Abnormal coupling of beta-adrenergic receptor to G protein is involved in the pathogenesis of the failing heart.

Selected Validation Data



Western blot analysis of GRK2 using anti-GRK2 antibody (A32388-1). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Human THP-1 whole cell lysates,

Lane 2: Monkey COS-7 whole cell lysates,


Lane 3: Human Raji whole cell lysates,

Lane 4: Rat lung tissue lysates,

Lane 5: Mouse NIH/3T3 whole cell lysates,


Lane 6: Mouse RAW264.7 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-GRK2 antigen affinity purified polyclonal antibody (A32388-1) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for GRK2 at approximately 80 kDa. The expected band size for GRK2 is at 80 kDa.



IF analysis of GRK2 using anti-GRK2 antibody (A32388-1).

GRK2 was detected in an immunocytochemical section of U2OS cells. The section was incubated with rabbit anti-GRK2 Antibody (A32388-1) at a dilution of 1:100. DyLight®488 Conjugated Goat Anti-Rabbit IgG (Green) (Catalog # BA1127) was used as secondary antibody. The section was counterstained with DAPI (Catalog # AR1176) (Blue).



Flow Cytometry analysis of Hepa1-6 cells using anti-GRK2 antibody (A32388-1).

Overlay histogram showing Hepa1-6 cells stained with A32388-1 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-GRK2 Antibody (A32388-1) at 1:100 dilution for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary

Product datasheet

Anti-GRK2 Antibody

Catalog Number: **A32388-1**



antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com

antibody and secondary antibody (Red line) was used as a blank control.