

## Basic Information

<b>Product Name</b>	Anti-Decorin/DCN Antibody		
<b>Gene Name</b>	DCN		
<b>Source</b>	Rabbit		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Species Reactivity</b>	human, mouse, rat, rabbit		
<b>Tested Application</b>	WB, IHC		
<b>Contents</b>	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.		
<b>Immunogen</b>	A synthetic peptide corresponding to a sequence at the C-terminus of human Decorin, identical to the related mouse sequence, and different from the related rat sequence by three amino acids.		
<b>Concentration</b>	500 ug/ml		
<b>Purification</b>	Immunogen affinity purified.		
<b>Observed MW</b>	40 kDa		
<b>Dilution Ratios</b>	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.		

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

## Background Information


Decorin is a proteoglycan on average 90-140 kilodaltons(kD) in size. The human decorin gene spans more than 38 kb and contains 8 exons and very large introns, 2 of which are 5.4 and more than 13.2 kb. And the human decorin gene is mapped to 12q21.3. It belongs to the small leucine-rich proteoglycan(SLRP) family and consists of a protein core containing leucine repeats with a glycosaminoglycan(GAG) chain consisting of either chondroitin sulfate(CS) or dermatan sulfate(DS). Decorin is a small cellular or pericellular matrix proteoglycan and is closely related in structure to biglycan protein. Decorin and biglycan are thought to be the result of a gene duplication. This protein is a component of connective tissue, binds to type I collagen fibrils, and plays a role in matrix assembly. And it also may play a role in epithelial/mesenchymal interactions during organ development and shaping. Decorin has been shown to have anti-

tumorigenic properties in an experimental murine tumor model and is capable of suppressing the growth of various tumor cell lines. There are multiple alternatively spliced transcript variants known for the decorin gene. Mutations in the decorin gene are associated with congenital stromal corneal dystrophy.

## Reference

Anti-Decorin/DCN Antibody被引用在3文献中。

## Selected Validation Data




Western blot analysis of Decorin/DCN using anti-Decorin/DCN antibody (BA0798). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Rat kidney tissue lysates,

Lane 2: Rat Spleen tissue lysates,

Lane 3: Rat Lung tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-Decorin/DCN antigen affinity purified polyclonal antibody (BA0798) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Decorin/DCN at approximately 40 kDa. The expected band size for Decorin/DCN is at 40 kDa.



IHC analysis of Decorin/DCN using anti-Decorin/DCN antibody (BA0798).

Decorin/DCN was detected in a paraffin-embedded section of human breast cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-Decorin/DCN Antibody (BA0798) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.

Product datasheet

## Anti-Decorin/DCN Antibody

Catalog Number: **BA0798**



antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,  
East Lake High-Tech Development Zone, Wuhan.

**Web:** [www.boster.com](http://www.boster.com) **Phone:** 027-67845390/1/2 **Email:** [boster@boster.com](mailto:boster@boster.com)