

Basic Information

Product Name	Anti-FGF4 Antibody
Gene Name	FGF4
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat, rabbit
Tested Application	IHC
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human FGF4, identical to the related rat and mouse sequences.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Dilution Ratios	Immunohistochemistry in paraffin section IHC-(P): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Fibroblast growth factor 4(FGF4), also known as Heparin Secretary Transforming(HSTF1). HST1, for which the designation HSTF1 was proposed for human gene nomenclature, is a heparin-binding growth factor with significant homology to human fibroblast growth factors and the mouse Int-2 protein. By in situ hybridization, Adelaide et al.(1988) mapped the HST gene to chromosome 11q13. The HST1 protein is a heparin-binding growth factor with significant homology with human fibroblast growth factors and the mouse Int-2 protein.

Selected Validation Data



IHC analysis of FGF4 using anti-FGF4 antibody (BA0848-1). FGF4 was detected in a paraffin-embedded section of human breast cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-FGF4 Antibody (BA0848-1) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.