BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

antibody and FLISA

Basic Information	
Product Name	Anti-VEGFB Antibody
Gene Name	VEGFB
Source	Rabbit
Clonality	Polyclonal
lsotype	lgG
Species Reactivity	human, mouse, rat, rabbit
Tested Application	IHC
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	E. coli-derived human VEGFB recombinant protein (Position: Q25-A138).
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Dilution Ratios	Immunohistochemistry in paraffin section IHC-(P): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Vascular endothelial growth factor B also known as VEGF-B is a protein that, in humans, is encoded by the VEGF-B gene. VEGF-B is a growth factor that belongs to the vascular endothelial growth factor family, of which VEGF-A is the best-known member. In contrast to VEGF-A, VEGF-B plays a less pronounced role in the vascular system: Whereas VEGF-A is important for the formation of blood vessels, such as during development or in pathological conditions, VEGF-B seems to play a role only in the maintenance of newly formed blood vessels during pathological conditions. It also plays an important role on several types of neurons. And it is important for the protection of neurons in the retina and thecerebral cortex during stroke and of motoneurons during motor neuron diseases such as amyotrophic lateral sclerosis.

Selected Validation Data

antibody and ELISA experts BOSTER BIOLOGICAL TECHNOLOGY Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com



IHC analysis of VEGFB using anti-VEGFB antibody (BA1460-1). VEGFB was detected in a paraffin-embedded section of rat cardiac muscle tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-VEGFB Antibody (BA1460-1) at a dilution of 1:200 and developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.