

## Basic Information

Product Name	Anti-ABCD3 Antibody	
Gene Name	ABCD3	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human PMP70, different from the related mouse and rat sequences by one amino acid.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	74 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	


## Storage

12 months from date of receipt, -20°C as supplied.

## Background Information

ATP-binding cassette sub-family D member 3 is a protein that in humans is encoded by the ABCD3 gene. The protein encoded by this gene is a member of the superfamily of ATP-binding cassette(ABC) transporters. The gene was assigned to human chromosome 1p21-p22 by in situ hybridization. This peroxisomal membrane protein likely plays an important role in peroxisome biogenesis. Mutations have been associated with some forms of Zellweger syndrome, a heterogeneous group of peroxisome assembly disorders.

## Selected Validation Data




Western blot analysis of ABCD3 using anti-ABCD3 antibody (BA3339). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Rat Lung tissue lysates,

Lane 2: Rat Ovary tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-ABCD3 antigen affinity purified polyclonal antibody (BA3339) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for ABCD3 at approximately 74 kDa. The expected band size for ABCD3 is at 75 kDa.



IHC analysis of ABCD3 using anti-ABCD3 antibody (BA3339).

ABCD3 was detected in a paraffin-embedded section of human lung cancer tissue. The tissue section was incubated with rabbit anti-ABCD3 Antibody (BA3339) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.