

Basic Information

Product Name	Anti-VCAM1 Antibody	
Gene Name	VCAM1	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	mouse, rat	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of mouse VCAM1, different from the related human sequence by three amino acids, and from the related rat sequence by two amino acids.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	85-110 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied.

Background Information


CD106(cluster of differentiation 106) also known as vascular cell adhesion molecule 1(VCAM-1), is a protein that in humans is encoded by the VCAM1 gene. In inflammatory conditions and in cardiac allografts undergoing rejection, VCAM1 is upregulated in endothelium of postcapillary venules. Arterial expression of VCAM1 is also found in experimental models of atherosclerosis in the rabbit. Cybulsky et al.(1991) mapped the VCAM1 gene to chromosome 1 by Southern analysis of somatic cell hybrids. Kumar et al.(1994) mapped the murine Vcam1 gene to chromosome 3 near Amy1. VCAM-1 functions as a cell adhesion molecule. The VCAM-1 protein mediates the adhesion of lymphocytes, monocytes, eosinophils, and basophils to vascular endothelium. It also functions in leukocyte-endothelial cell signal transduction, and it may play a role in the development of atherosclerosis and rheumatoid arthritis. CAM741 works similar to cotransin in that it represses the biosynthesis of VCAM1 cells by blocking the process of cotranslational translocation, which is dependent on the signal peptide of VCAM1. Among the lung metastasis

signature genes identified, several, including VCAM1, were functionally validated.

Reference

Anti-VCAM1 Antibody被引用在12文献中。

Selected Validation Data




Western blot analysis of VCAM1 using anti-VCAM1 antibody (BA3840). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Rat kidney tissue lysates,

Lane 2: Rat spleen tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-VCAM1 antigen affinity purified polyclonal antibody (BA3840) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for VCAM1 at approximately 85-110 kDa. The expected band size for VCAM1 is at 81 kDa.



IHC analysis of VCAM1 using anti-VCAM1 antibody (BA3840).

VCAM1 was detected in a paraffin-embedded section of rat intestinal tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-VCAM1 Antibody (BA3840) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.

Product datasheet

Anti-VCAM1 Antibody

Catalog Number: **BA3840**



antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com