

## Basic Information

Product Name	Anti-BAFF/TNFSF13B Antibody	
Gene Name	TNFSF13B	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human BAFF, different from the related rat and mouse sequences by three amino acids.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	31 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	


## Storage

12 months from date of receipt, -20°C as supplied.

## Background Information

BAFF was regularly detected by enzyme-linked immunosorbent assay in brain tissue lysates and in normal spinal fluid, and in astrocytes by double fluorescence microscopy. BAFF was localized in astrocytes close to BAFF-R-expressing immune cells. BAFF receptors were strongly expressed in situ in primary central nervous system (CNS) lymphomas. The TNF superfamily member B cell-activating factor (BAFF) plays an important role in humoral immunity and in autoimmune diseases, including RA. Local BAFF gene targeting inhibited proinflammatory cytokine expression, suppressed generation of plasma cells and Th17 cells, and markedly ameliorated joint pathology. The B cell activating factor BAFF (BlyS/TALL-1/zTNF4) is a tumor necrosis factor (TNF)-related ligand that promotes B cell survival and binds to three receptors (BCMA, TACI, and the recently described BAFF-R). Human BAFF was mapped to chromosome 13q32-34.4. The standard used in this kit is recombinant soluble human BAFF (A134-L295) with the molecular mass of 19.6 kDa.

## Selected Validation Data




Western blot analysis of BAFF/TNFSF13B using anti-BAFF/TNFSF13B antibody (BA3913).

Lane 1: recombinant human BAFF protein 10ng,

Lane 2: recombinant human BAFF protein 5ng,

Lane 3: recombinant human BAFF protein 2.5ng.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-BAFF/TNFSF13B antigen affinity purified polyclonal antibody (BA3913) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for BAFF/TNFSF13B at approximately 31 kDa.



IHC analysis of BAFF/TNFSF13B using anti-BAFF/TNFSF13B antibody (BA3913).

BAFF/TNFSF13B was detected in a paraffin-embedded section of human tonsil tissue. The tissue section was incubated with rabbit anti-BAFF/TNFSF13B Antibody (BA3913) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.

Product datasheet

## Anti-BAFF/TNFSF13B Antibody

Catalog Number: **BA3913**



antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,  
East Lake High-Tech Development Zone, Wuhan.

**Web:** [www.boster.com](http://www.boster.com) **Phone:** 027-67845390/1/2 **Email:** [boster@boster.com](mailto:boster@boster.com)