

## Basic Information

|                           |   |            |
|---------------------------|---|------------|
| <b>Product Name</b>       | Anti-active Caspase 3/CASP3 (p17) Antibody (Clone#HE-3)   |            |
| <b>Gene Name</b>          | CASP3   |            |
| <b>Source</b>             | Rabbit  |            |
| <b>Clonality</b>          | Monoclonal  |            |
| <b>Isotype</b>            | IgG   |            |
| <b>Species Reactivity</b> | human   |            |
| <b>Tested Application</b> | WB,IHC,ICC/IF   |            |
| <b>Contents</b>           | 500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol. |            |
| <b>Immunogen</b>          | A synthesized peptide derived from human active Caspase-3   |            |
| <b>Concentration</b>      | 500 ug/ml   |            |
| <b>Purification</b>       | Affinity-chromatography   |            |
| <b>Observed MW</b>        | 35 kDa,(cleaved)20/17/12 kDa  |            |
| <b>Dilution Ratios</b>    | Western blot (WB):  | 1:500-2000 |
|                           | Immunohistochemistry (IHC):   | 1:50-200   |
|                           | Immunocytochemistry/Immunofluorescence (ICC/IF):  | 1:50-200   |

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

## Background Information

Caspase 3 is a caspase protein which interacts with Survivin, XIAP, CFLAR, Caspase 8, HCLS1, Deleted in Colorectal Cancer, TRAF3 and GroEL. This gene which is located on 4q35 encodes a protein that is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes that undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. And the caspase-3 activation in heart failure sequentially cleaves SRF and generates a truncated SRF that appears to function as a dominant-negative transcription factor. Additionally, the caspase-3 influence on bone mineral density should be considered in any in vivo application of caspase-3 inhibitors to the treatment of human disease. In erythroid precursors undergoing terminal differentiation, Hsp70

Product datasheet

## Anti-active Caspase 3/CASP3 (p17)

Antibody (Clone#HE-3)

Catalog Number: **BM3937**

# BOSTER

antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator,  
Wuhan East Lake High-tech Development Zone


Web: [www.boster.com](http://www.boster.com) Phone: 027-67845390 Email: [boster@boster.com](mailto:boster@boster.com)

prevents active CASP3 from cleaving GATA1 and inducing apoptosis.

## Reference

Anti-active Caspase 3/CASP3 (p17) Antibody (Clone#HE-3)被引用在15文献中。

## Selected Validation Data



Western blot analysis of active Caspase-3 expression in Jurkat cell lysate treated with Camptothecin.

Product datasheet

## Anti-active Caspase 3/CASP3 (p17)

Antibody (Clone#HE-3)

Catalog Number: **BM3937**


# BOSTER

antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator,  
Wuhan East Lake High-tech Development Zone

**Web:** [www.boster.com](http://www.boster.com) **Phone:** 027-67845390 **Email:** [boster@boster.com](mailto:boster@boster.com)



Immunofluorescent analysis of HepG2 cells, using active Caspase-3 Antibody .