

Basic Information

| | | |
|---------------------------|---|------------|
| Product Name | Anti-pro Caspase 3/CASP3 Antibody (Clone#AOB-3) | |
| Gene Name | CASP3 | |
| Source | Rabbit | |
| Clonality | Monoclonal | |
| Isotype | IgG | |
| Species Reactivity | human, mouse | |
| Tested Application | WB, IHC, ICC/IF, IP, FCM | |
| Contents | 500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol. | |
| Immunogen | A synthesized peptide derived from human pro Caspase 3 | |
| Concentration | 500 ug/ml | |
| Purification | Affinity-chromatography | |
| Observed MW | 35 kDa,(cleaved)20/17/12 kDa | |
| Dilution Ratios | Western blot (WB): | 1:500-2000 |
| | Immunohistochemistry (IHC): | 1:50-200 |
| | Immunocytochemistry/Immunofluorescence (ICC/IF): | 1:50-200 |
| | ImmunoPrecipitation (IP): | 1:50 |
| | Flow Cytometry (FCM): | 1:50 |

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information


Caspase 3 is a caspase protein which interacts with Survivin, XIAP, CFLAR, Caspase 8, HCLS1, Deleted in Colorectal Cancer, TRAF3 and GroEL. This gene which is located on 4q35 encodes a protein that is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes that undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. And the caspase-3 activation in heart failure sequentially cleaves SRF and generates a truncated SRF that appears to function as a dominant-negative transcription factor. Additionally, the caspase-3 influence on bone mineral density should be considered in any in vivo application of caspase-3

inhibitors to the treatment of human disease. In erythroid precursors undergoing terminal differentiation, Hsp70 prevents active CASP3 from cleaving GATA1 and inducing apoptosis.

Reference

Anti-pro Caspase 3/CASP3 Antibody (Clone#AOB-3)被引用在27文献中。

Selected Validation Data



Western blot analysis of pro Caspase 3 in HeLa cell lysate.

Product datasheet

**Anti-pro Caspase 3/CASP3 Antibody
(Clone#AOB-3)**

Catalog Number: BM3954

BOSTER


antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd and 4th floors, Optics Valley Biomedical Accelerator,
Wuhan East Lake High-tech Development Zone

Web: www.boster.com **Phone:** 027-67845390 **Email:** boster@boster.com

pro Caspase 3 Antibody



Immunofluorescent analysis of A673 cells, using pro Caspase 3 Antibody.