

Basic Information

Product Name	Anti-CDC37 Antibody (Clone#ICA-3)		
Gene Name	CDC37		
Source	Rabbit		
Clonality	Monoclonal		
Isotype	IgG		
Species Reactivity	human, mouse, rat		
Tested Application	WB, IHC, IP, FCM		
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.		
Immunogen	A synthesized peptide derived from human Cdc37		
Concentration	500 ug/ml		
Purification	Affinity-chromatography		
Observed MW	44 kDa		
Dilution Ratios	Western blot (WB):	1:500-2000	
	Immunohistochemistry (IHC):	1:50-200	
	ImmunoPrecipitation (IP):	1:20	
	Flow Cytometry (FCM):	1:20	


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Hsp90 co-chaperone Cdc37, also called P50CDC37, is a protein that in humans is encoded by the CDC37 gene. The protein encoded by this gene is highly similar to Cdc 37, a cell division cycle control protein of *Saccharomyces cerevisiae*. This gene is mapped to 19p13.2. And this protein is a molecular chaperone with specific function in cell signal transduction. It has been shown to form complex with Hsp90 and a variety of protein kinases including CDK4, CDK6, SRC, RAF-1, MOK, as well as eIF2 alpha kinases. It is thought to play a critical role in directing Hsp90 to its target kinases.

Selected Validation Data



Western blot analysis of Cdc37 expression in (1) Jurkat cell lysate; (2) SW480 cell lysate.