

Basic Information

Product Name	Anti-CYP19A1 Antibody (Clone#AABB-3)
Gene Name	CYP19A1
Source	Rabbit
Clonality	Monoclonal
Isotype	IgG
Species Reactivity	human, rat
Tested Application	WB, IP
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.
Immunogen	A synthesized peptide derived from human Aromatase
Concentration	500 ug/ml
Purification	Affinity-chromatography
Observed MW	58 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 ImmunoPrecipitation (IP):1:20

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Aromatase is an enzyme responsible for a key step in the biosynthesis of estrogens. It is a member of the cytochrome P450 superfamily, which are monooxygenases that catalyze many reactions involved in steroidogenesis. In particular, aromatase is responsible for the aromatization of androgens into estrogens. The CYP19 gene spans at least 70 kb of genomic DNA and contains 10 exons. By in situ hybridization, the ARO gene is mapped to 15q21.1. The aromatase enzyme can be found in many tissues including gonads, brain, adipose tissue, placenta, blood vessels, skin, bone, and endometrium, as well as in tissue of endometriosis, uterine fibroids, breast cancer, and endometrial cancer. It is an important factor in sexual development. Some bodybuilders taking steroids also take antiaromatase supplements to prevent excess testosterone conversion into estrogens, which can cause gynecomastia.

Reference

Anti-CYP19A1 Antibody (Clone#AABB-3)被引用在2文献中。

Selected Validation Data

Western blot analysis of Aromatase expression in rat brain lysate.

