

Basic Information

Product Name	Anti-NR1D1 Antibody (Clone#18N41)	
Gene Name	NR1D1	
Source	Rabbit	
Clonality	Monoclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF, FCM	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
Immunogen	A synthesized peptide derived from human NR1D1	
Concentration	500 ug/ml	
Observed MW	55 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-200
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-200
	Flow Cytometry (FCM):	1:20

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Functions as a constitutive transcriptional repressor. Possible receptor for triiodothyronine.

Reference

Anti-NR1D1 Antibody (Clone#18N41)被引用在1文献中。

Selected Validation Data



Western blot analysis of anti-NR1D1 antibody (BM5531). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human HepG2 whole cell lysates,

Lane 3: human U-87MG whole cell lysates,

Lane 4: human U2OS whole cell lysates,

Lane 5: rat liver tissue lysates,

Lane 6: rat RH-35 whole cell lysates,

Lane 7: mouse liver tissue lysates,

Lane 8: mouse HEPA1-6 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-NR1D1 antigen affinity purified monoclonal antibody (BM5531) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for NR1D1 at approximately 55 kDa. The expected band size for NR1D1 is at 67 kDa.