

Basic Information

| | |
|---------------------------|--|
| Product Name | Anti-P glycoprotein/ABCB1 Antibody (Clone#OTI2G4) |
| Gene Name | ABCB1 |
| Source | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 |
| Species Reactivity | human, mouse, rat |
| Tested Application | WB |
| Contents | PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Immunogen | Human recombinant protein fragment corresponding to amino acids 347-710 of human ABCB1 (NP_000918) produced in SF9 cell. |
| Concentration | 500 ug/ml |
| Purification | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Dilution Ratios | Western blot (WB):1:2000 |


Storage

Stable for 12 months from date of receipt. Store at -20°C as received.

Background Information

P-GP, also called ABCB1 or PGY1, is a glycoprotein that in humans is encoded by the ABCB1 gene. It is mapped to 7q21.12. P-GP is a well-characterized ABC-transporter (which transports a wide variety of substrates across extra- and intracellular membranes) of the MDR/TAP subfamily. It is an important protein of the cell membrane that pumps many foreign substances out of cells. More formally, it is an ATP-dependent drug efflux pump with broad substrate specificity. P-GP is an ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier.

Selected Validation Data



SF9 cells lysate (5 ug, left lane) and SF9 cells lysate expressing human recombinant protein fragment (5 ug, right lane) corresponding to amino acids 995-1280 of human ABCB1 were separated by SDS-PAGE and immunoblotted with anti-ABCB1.