


## Basic Information

<b>Product Name</b>	Anti-HNF4A Antibody (Clone#28H20)	
<b>Gene Name</b>	HNF4A	
<b>Source</b>	Rabbit	
<b>Clonality</b>	Monoclonal	
<b>Isotype</b>	IgG	
<b>Species Reactivity</b>	human, mouse, rat	
<b>Tested Application</b>	WB, IHC, ICC/IF, FCM	
<b>Contents</b>	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
<b>Immunogen</b>	A synthesized peptide derived from human HNF-4-alpha	
<b>Concentration</b>	500 ug/ml	
<b>Purification</b>	Affinity-chromatography	
<b>Observed MW</b>	53 kDa	
<b>Dilution Ratios</b>	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-200
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-200
	Flow Cytometry (FCM):	1:50

## Storage

12 months from date of receipt, -20°C as supplied.

## Selected Validation Data



Western blot analysis of anti-HNF4A antibody (M00389-4). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Caco-2 whole cell lysates,

Lane 2: human A549 whole cell lysates,

Lane 3: human Hela whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-HNF4A antigen

affinity purified monoclonal antibody (M00389-4) at a dilution of

1:1000 and probed with a goat anti-rabbit IgG-HRP secondary

antibody (Catalog # BA1054). The signal is developed using ECL Plus

Western Blotting Substrate (Catalog # AR1197). A specific band was

detected for HNF4A at approximately 53 kDa. The expected band

size for HNF4A is at 53 kDa.