

Basic Information

Product Name	Anti-IGF2R Antibody (Clone#6G2)	
Gene Name	IGF2R	
Source	Mouse	
Clonality	Monoclonal	
Isotype	IgG2b	
Species Reactivity	human	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E. coli-derived human IGF2R recombinant protein (Position: F424-R529).	
Concentration	500 ug/ml	
Purification	protein G purified.	
Observed MW	274 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry in paraffin section IHC 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer, pH6.0, or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Insulin-like growth factor 2 receptor, also called IGF2R or I-MPR is a protein that in humans is encoded by the IGF2R gene. This gene is mapped to 6q25.3. This gene encodes a receptor for both insulin-like growth factor 2 and mannose 6-phosphate, although the binding sites for either are located on different segments of the receptor. This receptor functions in the intracellular trafficking of lysosomal enzymes, the activation of transforming growth factor beta, and the degradation of insulin-like growth factor 2. While the related mouse gene shows exclusive expression from the maternal allele, imprinting of the human gene appears to be polymorphic, with only a minority of individuals showing expression from the maternal allele.

Selected Validation Data




Western blot analysis of IGF2R using anti-IGF2R antibody (M00951-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,

Lane 2: human U87 whole cell lysates,

Lane 3: human HEPG2 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with mouse anti-IGF2R antigen affinity purified monoclonal antibody (M00951-2) at a dilution of 1:1000 and probed with a goat anti-mouse IgG-HRP secondary antibody (Catalog # BA1050). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for IGF2R at approximately 274 kDa. The expected band size for IGF2R is at 274 kDa.



IHC analysis of IGF2R using anti-IGF2R antibody (M00951-2).

IGF2R was detected in a paraffin-embedded section of human placenta tissue. Biotinylated goat anti-mouse IgG was used as secondary antibody. The tissue section was incubated with mouse anti-IGF2R Antibody (M00951-2) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB (Catalog # AR1027) as the chromogen.