

Basic Information

| | | |
|---------------------------|--|------------|
| Product Name | Anti-BubR1/BUB1B Antibody (Clone#8B3) | |
| Gene Name | BUB1B | |
| Source | Mouse | |
| Clonality | Monoclonal | |
| Isotype | IgG2b | |
| Species Reactivity | human, rat | |
| Tested Application | WB, IHC, FCM | |
| Contents | 500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol. | |
| Immunogen | E.coli-derived human BubR1/BUB1B recombinant protein (Position: K26-E448). | |
| Concentration | 500 ug/ml | |
| Purification | protein G purified. | |
| Observed MW | 130 kDa | |
| Dilution Ratios | Western blot (WB): | 1:500-2000 |
| | Immunohistochemistry (IHC): | 1:50-400 |
| | Flow Cytometry (Fixed): | 1:50-200 |
| | (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user. | |


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Mitotic checkpoint serine/threonine-protein kinase BUB1 beta is an enzyme that in humans is encoded by the BUB1B gene. This gene encodes a kinase involved in spindle checkpoint function. The protein has been localized to the kinetochore and plays a role in the inhibition of the anaphase-promoting complex/cyclosome (APC/C), delaying the onset of anaphase and ensuring proper chromosome segregation. Impaired spindle checkpoint function has been found in many forms of cancer.

Selected Validation Data



Western blot analysis of BubR1/BUB1B using anti-BubR1/BUB1B antibody (M01564-3). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.


Lane 1: human HeLa whole cell lysates,

Lane 2: human HEK293 whole cell lysates,

Lane 3: human K562 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with mouse anti-BubR1/BUB1B antigen affinity purified monoclonal antibody (M01564-3) at a dilution of 1:1000 and probed with a goat anti-mouse IgG-HRP secondary antibody (Catalog # BA1050). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for BubR1/BUB1B at approximately 130 kDa. The expected band size for BubR1/BUB1B is at 120 kDa.



IHC analysis of BubR1/BUB1B using anti-BubR1/BUB1B antibody (M01564-3).

BubR1/BUB1B was detected in a paraffin-embedded section of human mammary cancer tissue. Biotinylated goat anti-mouse IgG was used as secondary antibody. The tissue section was incubated with mouse anti-BubR1/BUB1B Antibody (M01564-3) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB (Catalog # AR1027) as the chromogen.

Product datasheet
Anti-BubR1/BUB1B Antibody
(Clone#8B3)
Catalog Number: M01564-3


BOSTER[®]

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com



Flow Cytometry analysis of Hela cells using anti-BubR1/BUB1B antibody (M01564-3).

Overlay histogram showing Hela cells stained with M01564-3 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-BubR1/BUB1B Antibody (M01564-3) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.